


## CLAIMS

1. A compound of formula (I),



the *N*-oxide forms, the addition salts and the stereo-chemically isomeric forms thereof, wherein

10 *n* is 0 or 1;  
*s* is 0 or 1;

*X* is -N= or -CR<sup>4</sup>=, wherein R<sup>4</sup> is hydrogen or taken together with R<sup>1</sup> may form a bivalent radical of formula -CH=CH-CH=CH-;

15 *Y* is -N< or -CH<;

*Q* is -NH-, -O-, -C(O)-, -CH<sub>2</sub>-CH<sub>2</sub>- or -CHR<sup>5</sup>-,  
 wherein R<sup>5</sup> is hydrogen, hydroxy, C<sub>1-6</sub>alkyl, arylC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxycarbonyl,  
 20 C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkylamino or haloindazolyl;

R<sup>1</sup> is C<sub>1-6</sub>alkyl or thienyl;

R<sup>2</sup> is hydrogen or taken together with R<sup>3</sup> may form =O;

25 R<sup>3</sup> is hydrogen, C<sub>1-6</sub>alkyl or a radical selected from

- NR<sup>6</sup>R<sup>7</sup> (a-1),
- O-H (a-2),
- O-R<sup>8</sup> (a-3),
- 30 - S- R<sup>9</sup> (a-4), or
- C≡N (a-5),

wherein

R<sup>6</sup> is -CHO, C<sub>1-6</sub>alkyl, hydroxyC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonyl,  
 di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonylaminoC<sub>1-6</sub>alkyl,

piperidinylC<sub>1-6</sub>alkyl, piperidinylC<sub>1-6</sub>alkylaminocarbonyl, C<sub>1-6</sub>alkyloxy,  
 C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl, thienylC<sub>1-6</sub>alkyl, pyrrolylC<sub>1-6</sub>alkyl,  
 arylC<sub>1-6</sub>alkylpiperidinyl, arylcarbonylC<sub>1-6</sub>alkyl, arylcarbonylpiperidinylC<sub>1-6</sub>alkyl,  
 haloindazolylpiperidinylC<sub>1-6</sub>alkyl, or arylC<sub>1-6</sub>alkyl(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl; and  
 5 R<sup>7</sup> is hydrogen or C<sub>1-6</sub>alkyl;  
 R<sup>8</sup> is C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonyl or di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl; and  
 R<sup>9</sup> is di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl;














or R<sup>3</sup> is a group of formula




wherein

t is 0, 1 or 2;

Z is a heterocyclic ring system selected from



wherein each R<sup>10</sup> independently is hydrogen, C<sub>1-6</sub>alkyl, aminocarbonyl, hydroxy,




C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkylamino, di(phenylC<sub>2-6</sub>alkenyl),  
 piperidinylC<sub>1-6</sub>alkyl, C<sub>3-10</sub>cycloalkyl, C<sub>3-10</sub>cycloalkylC<sub>1-6</sub>alkyl,  
 aryloxy(hydroxy)C<sub>1-6</sub>alkyl, haloindazolyl, arylC<sub>1-6</sub>alkyl, arylC<sub>2-6</sub>alkenyl,  
 25 morpholino, C<sub>1-6</sub>alkylimidazolyl, or pyridinylC<sub>1-6</sub>alkylamino;

each R<sup>11</sup> independently is hydrogen, hydroxy, piperidinyl or aryl;


aryl is phenyl or phenyl substituted with halo, C<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxy;

- 5 with the proviso that 6-(cyclohexyl-1*H*-imidazol-1-ylmethyl)-3-methyl-2(1*H*)-quinoxalinone is not included.
2. A compound as claimed in claim 1 wherein X is -N= or -CH=; R<sup>1</sup> is C<sub>1-6</sub>alkyl; R<sup>3</sup> is hydrogen, C<sub>1-6</sub>alkyl, a radical selected from (a-1), (a-2), (a-3) or (a-4) or a group of  
 10 formula (b-1); R<sup>6</sup> is di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl or C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl; R<sup>7</sup> is hydrogen; R<sup>8</sup> is di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl; t is 0 or 2; Z is a heterocyclic ring system selected from (c-1), (c-5), (c-6), (c-8), (c-10), (c-12) or (c-13); each R<sup>10</sup> independently is hydrogen, C<sub>1-6</sub>alkyl, hydroxy, C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkylamino, morpholino, C<sub>1-6</sub>alkylimidazolyl, or  
 15 pyridinylC<sub>1-6</sub>alkylamino; each R<sup>11</sup> independently is hydrogen or hydroxy; and aryl is phenyl.
3. A compound according to claim 1 and 2 wherein n is 0; X is CH; Q is -NH-, -CH<sub>2</sub>-CH<sub>2</sub>- or -CHR<sup>5</sup>-, wherein R<sup>5</sup> is hydrogen, hydroxy, or arylC<sub>1-6</sub>alkyl; R<sup>1</sup> is C<sub>1-6</sub>alkyl; R<sup>2</sup> is hydrogen; R<sup>3</sup> is hydrogen, hydroxy or a group of formula (b-1); t is 0; Z is a heterocyclic ring system selected from (c-8) or (c-13); each R<sup>10</sup> independently is hydrogen; and aryl is phenyl.
4. A compound according to claim 1, 2 and 3 wherein the compound is selected from  
 25 compound No 7, compound No 2, compound No 1 and compound No 11.



5. A compound as claimed in any of claims 1 to 4 for use as a medicine.

6. A pharmaceutical composition comprising pharmaceutically acceptable carriers and as an active ingredient a therapeutically effective amount of a compound as claimed in claim 1 to 4.
- 5 7. A process of preparing a pharmaceutical composition as claimed in claim 6 wherein the pharmaceutically acceptable carriers and a compound as claimed in claim 1 to 4 are intimately mixed.
8. Use of a compound for the manufacture of a medicament for the treatment of a
- 10 PARP mediated disorder, wherein the compound is a compound of formula (I)



- 15 the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereochemically isomeric forms thereof, wherein

*n* is 0 or 1;

*s* is 0 or 1;

20

*X* is  $-N=$  or  $-CR^4=$ , wherein  $R^4$  is hydrogen or taken together with  $R^1$  may form a bivalent radical of formula  $-CH=CH-CH=CH-$ ;

*Y* is  $-N<$  or  $-CH<$ ;

25

*Q* is  $-NH-$ ,  $-O-$ ,  $-C(O)-$ ,  $-CH_2-CH_2-$  or  $-CHR^5-$ ,  
wherein  $R^5$  is hydrogen, hydroxy,  $C_{1-6}$ alkyl, aryl $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxycarbonyl,  $C_{1-6}$ alkyloxy $C_{1-6}$ alkylamino or haloindazolyl;

30  $R^1$  is  $C_{1-6}$ alkyl or thienyl;

$R^2$  is hydrogen or taken together with  $R^3$  may form  $=O$ ;

$R^3$  is hydrogen,  $C_{1-6}$ alkyl or a radical selected from

- 5      - NR<sup>6</sup>R<sup>7</sup>                      (a-1),  
          -O-H                      (a-2),  
          -O-R<sup>8</sup>                      (a-3),  
          -S- R<sup>9</sup>                      (a-4), or  
          —C≡N                      (a-5),

wherein

- R<sup>6</sup> is -CHO, C<sub>1-6</sub>alkyl, hydroxyC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonyl,  
 di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonylaminoC<sub>1-6</sub>alkyl,  
 piperidinyC<sub>1-6</sub>alkyl, piperidinyC<sub>1-6</sub>alkylaminocarbonyl, C<sub>1-6</sub>alkyloxy,  
 10 C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl, thienylC<sub>1-6</sub>alkyl, pyrrolylC<sub>1-6</sub>alkyl,  
 arylC<sub>1-6</sub>alkylpiperidiny, arylcarbonylC<sub>1-6</sub>alkyl, arylcarbonylpiperidinyC<sub>1-6</sub>alkyl,  
 haloindozolylpiperidinyC<sub>1-6</sub>alkyl, or arylC<sub>1-6</sub>alkyl(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl; and  
 R<sup>7</sup> is hydrogen or C<sub>1-6</sub>alkyl;  
 R<sup>8</sup> is C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonyl or di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl; and  
 15 R<sup>9</sup> is di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl;














or R<sup>3</sup> is a group of formula



wherein


t is 0, 1 or 2;

- 20      Z is a heterocyclic ring system selected from



-53-

wherein each  $R^{10}$  independently is hydrogen,  $C_{1-6}$ alkyl, aminocarbonyl, hydroxy,



$C_{1-6}$ alkyloxy $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy $C_{1-6}$ alkylamino, di(phenyl $C_{2-6}$ alkenyl),  
 piperidinyl $C_{1-6}$ alkyl,  $C_{3-10}$ cycloalkyl,  $C_{3-10}$ cycloalkyl $C_{1-6}$ alkyl,  
 5 aryloxy(hydroxy) $C_{1-6}$ alkyl, haloindazolyl, aryl $C_{1-6}$ alkyl, aryl $C_{2-6}$ alkenyl,  
 morpholino,  $C_{1-6}$ alkylimidazolyl, or pyridinyl $C_{1-6}$ alkylamino;  
 each  $R^{11}$  independently is hydrogen, hydroxy, piperidinyl or aryl;


aryl is phenyl or phenyl substituted with halo,  $C_{1-6}$ alkyl, or  $C_{1-6}$ alkyloxy.

9. Use according to claim 8 of a PARP inhibitor of formula (I) for the manufacture of  
 a medicament for the treatment of a PARP-1 mediated disorder

10. Use according to claim 8 and 9 wherein the treatment involves chemosensitization.

11. Use according to claims 8 and 9 wherein the treatment involves radiosensitization.

12. A combination of a compound with a chemotherapeutic agent wherein said  
 compound is a compound of formula (I)



the *N*-oxide forms, the pharmaceutically acceptable addition salts and the stereo-  
 chemically isomeric forms thereof, wherein

$n$  is 0 or 1;

$s$  is 0 or 1;

$X$  is  $-N=$  or  $-CR^4=$ , wherein  $R^4$  is hydrogen or taken together with  $R^1$  may form a  
 bivalent radical of formula  $-CH=CH-CH=CH-$ ;

Y is  $-N<$  or  $-CH<$ ;

Q is  $-NH-$ ,  $-O-$ ,  $-C(O)-$ ,  $-CH_2-CH_2-$  or  $-CHR^5-$ ,

wherein  $R^5$  is hydrogen, hydroxy,  $C_{1-6}$ alkyl, aryl $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxycarbonyl,

5  $C_{1-6}$ alkyloxy $C_{1-6}$ alkylamino or haloindazolyl;

$R^1$  is  $C_{1-6}$ alkyl or thienyl;

$R^2$  is hydrogen or taken together with  $R^3$  may form  $=O$ ;

10

$R^3$  is hydrogen,  $C_{1-6}$ alkyl or a radical selected from

$-NR^6R^7$  (a-1),

$-O-H$  (a-2),

$-O-R^8$  (a-3),

15  $-S-R^9$  (a-4), or

$-C\equiv N$  (a-5),

wherein

$R^6$  is  $-CHO$ ,  $C_{1-6}$ alkyl, hydroxy $C_{1-6}$ alkyl,  $C_{1-6}$ alkylcarbonyl,

di( $C_{1-6}$ alkyl)amino $C_{1-6}$ alkyl,  $C_{1-6}$ alkylcarbonylamino $C_{1-6}$ alkyl,

20 piperidinyl $C_{1-6}$ alkyl, piperidinyl $C_{1-6}$ alkylaminocarbonyl,  $C_{1-6}$ alkyloxy,

$C_{1-6}$ alkyloxy $C_{1-6}$ alkyl, thienyl $C_{1-6}$ alkyl, pyrrolyl $C_{1-6}$ alkyl,

aryl $C_{1-6}$ alkylpiperidinyl, arylcarbonyl $C_{1-6}$ alkyl, arylcarbonylpiperidinyl $C_{1-6}$ alkyl,

haloindazolylpiperidinyl $C_{1-6}$ alkyl, or aryl $C_{1-6}$ alkyl( $C_{1-6}$ alkyl)amino $C_{1-6}$ alkyl; and

$R^7$  is hydrogen or  $C_{1-6}$ alkyl;

25  $R^8$  is  $C_{1-6}$ alkyl,  $C_{1-6}$ alkylcarbonyl or di( $C_{1-6}$ alkyl)amino $C_{1-6}$ alkyl; and

$R^9$  is di( $C_{1-6}$ alkyl)amino $C_{1-6}$ alkyl;

or  $R^3$  is a group of formula


$-(CH_2)_t-Z-$  (b-1),

wherein

30

t is 0, 1 or 2;


Z is a heterocyclic ring system selected from




(c-1)



(c-2)




(c-3)




(c-4)


-55-




(c-5)




(c-6)




(c-7)




(c-8)




(c-9)




(c-10)



(c-11)




(c-12)



(c-13)

5 wherein each  $R^{10}$  independently is hydrogen,  $C_{1-6}$ alkyl, aminocarbonyl, hydroxy,




$C_{1-6}$ alkyloxy $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy $C_{1-6}$ alkylamino, di(phenyl $C_{2-6}$ alkenyl),  
 piperidinyl $C_{1-6}$ alkyl,  $C_{3-10}$ cycloalkyl,  $C_{3-10}$ cycloalkyl $C_{1-6}$ alkyl,  
 aryloxy(hydroxy) $C_{1-6}$ alkyl, haloindazolyl, aryl $C_{1-6}$ alkyl, aryl $C_{2-6}$ alkenyl,  
 10 morpholino,  $C_{1-6}$ alkylimidazolyl, or pyridinyl $C_{1-6}$ alkylamino;  
 each  $R^{11}$  independently is hydrogen, hydroxy, piperidinyl or aryl;

aryl is phenyl or phenyl substituted with halo,  $C_{1-6}$ alkyl or  $C_{1-6}$ alkyloxy.


- 15 13. A process for preparing a compound as claimed in claim 1, characterized by  
 a) the hydrolysis of intermediates of formula (VIII), according to art-known methods,  
 by submitting the intermediates of formula (VIII) to appropriate reagents, such as,  
 tinchloride, acetic acid and hydrochloric acid, in the presence of a reaction inert  
 solvent, e.g. tetrahydrofuran.

20






- b) the cyclization of intermediates of formula (X), according to art-known cyclizing procedures into compounds of formula (I) wherein X is CH herein referred to as compounds of formula (I-j), preferably in the presence of a suitable Lewis Acid, e.g. aluminum chloride either neat or in a suitable solvent such as, for example, an aromatic hydrocarbon, e.g. benzene, chlorobenzene, methylbenzene and the like; halogenated hydrocarbons, e.g. trichloromethane, tetrachloromethane and the like; an ether, e.g. tetrahydrofuran, 1,4-dioxane and the like or mixtures of such solvents.



10

- c) the condensation of an appropriate ortho-benzenediamine of formula (XI) with an ester of formula (XII) into compounds of formula (I), wherein X is N and R<sup>2</sup> taken together with R<sup>3</sup> forms =O, herein referred to as compounds of formula (I-a-1), in the presence of a carboxylic acid, e.g. acetic acid and the like, a mineral acid such as, for example hydrochloric acid, sulfuric acid, or a sulfonic acid such as, for example, methanesulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid and the like.



20